

INTRODUCTION TO HECKE ALGEBRAS

ELAD ZELINGHER

Let F be a non-archimedean local field with ring of integers \mathfrak{o} , maximal ideal \mathfrak{p} , uniformizer $\varpi \in \mathfrak{p} \setminus \mathfrak{o}$ and residue field \mathbb{F}_q . Let G be a split connected algebraic reductive group over F , with split maximal torus A and Borel subgroup $B = AN$. Let $\mathcal{K} = G(\mathfrak{o})$ be a hyperspecial maximal open compact subgroup of G . Let W be the Weyl group of G , that is, the normalizer of A in G modulo A .

Let $X_*(A)$ denote the cocharacter group

$$X_*(A) = \text{Hom}(\mathbb{G}_m, A) = \{ \mu^\vee : F^\times \rightarrow A \mid \mu^\vee \text{ algebraic character} \}.$$

For an element $\mu^\vee \in X_*(A)$ we denote $\varpi^{\mu^\vee} := \mu^\vee(\varpi)$. The map $\mu \mapsto \varpi^{\mu^\vee}$ defines an isomorphism

$$X_*(A) \rightarrow A(\varpi) := A/A(\mathfrak{o}),$$

where $A(\mathfrak{o}) = A \cap \mathcal{K}$.

Example 1. When $G = \text{GL}_n$,

$$A = \left\{ \begin{pmatrix} a_1 & & \\ & \ddots & \\ & & a_n \end{pmatrix} \mid a_1, \dots, a_n \in F^\times \right\}$$

and

$$X_*(A) \cong \mathbb{Z}^n$$

by the isomorphism sending

$$\mathbb{Z} \ni (k_1, \dots, k_n) \mapsto \mu_{(k_1, \dots, k_n)}^\vee(x) = \begin{pmatrix} x^{k_1} & & \\ & \ddots & \\ & & x^{k_n} \end{pmatrix}.$$

In this case,

$$A(\varpi) = A/A(\mathfrak{o}) \cong \left\{ \begin{pmatrix} \varpi^{k_1} & & \\ & \ddots & \\ & & \varpi^{k_n} \end{pmatrix} \mid k_1, \dots, k_n \in \mathbb{Z} \right\},$$

where

$$A(\mathfrak{o}) = \left\{ \begin{pmatrix} a_1 & & \\ & \ddots & \\ & & a_n \end{pmatrix} \mid a_1, \dots, a_n \in \mathfrak{o}^\times \right\}.$$

It is clear that the map $X_*(A) \rightarrow A(\varpi)$ given by $(k_1, \dots, k_n) \mapsto \mu_{(k_1, \dots, k_n)}^\vee(\varpi)$ is an isomorphism.

For any compact open subgroup $K \subset G$ let

$$\mathcal{H}(G // K) = \{f \in C_c^\infty(G) \mid f(kgk') = f(g), \forall g \in G, k, k' \in K\}.$$

A basis for $\mathcal{H}(G // K)$ is given by the characteristic functions $\{\chi_{KgK}\}_g$ where g runs over all the representatives of cosets of $K \backslash G / K$.

The space $\mathcal{H}(G // K)$ is an algebra with convolution as multiplication:

$$(f_1 * f_2)(g) = \int_G f_1(x^{-1}) f_2(xg) dx.$$

1. SPHERICAL HECKE ALGEBRA

Recall the Cartan decomposition: we have that

$$G = \bigcup_{\substack{\mu^\vee \\ \mu^\vee \text{ is dominant}}} \mathcal{K} \cdot \varpi^{\mu^\vee} \cdot \mathcal{K}.$$

It follows from applying Gelfand's trick to an involution based on the Chevalley data, that $\mathcal{H}(G // \mathcal{K})$ is commutative (the involution should act trivially on ϖ^{μ^\vee} for any μ^\vee and send \mathcal{K} to itself).

Example 2. If $G = \mathrm{GL}_n(F)$, then $\mu_{(k_1, \dots, k_n)}^\vee$ is dominant if and only if $k_1 \geq k_2 \geq \dots \geq k_n$. The involution in this case is $x \mapsto {}^t x$. In this case, under the isomorphism above, every dominant cocharacter can be written as a product of the form

$$a_1^{i_1} \cdot a_2^{i_2} \cdot \dots \cdot a_n^{i_n},$$

where $i_1, \dots, i_n \in \mathbb{Z}$ and $i_1, \dots, i_{n-1} \geq 0$, and for $1 \leq j \leq n$,

$$a_j = \begin{pmatrix} \varpi I_j & \\ & I_{n-j} \end{pmatrix}.$$

The spherical Hecke algebra $\mathcal{H}(G // \mathcal{K})$ can be realized as compactly supported \mathbb{C} -valued functions on $A(\varpi)$, that are invariant under the action of the Weyl group W .

If $K \subset \mathcal{K}$ is an arbitrary compact open subgroup, then $\mathcal{H}(G // K)$ is not commutative when $K \neq \mathcal{K}$. However, we have the following relation. If μ_1^\vee and μ_2^\vee are dominant cocharacters then

$$\chi_{K \cdot \varpi^{\mu_1^\vee} \cdot K} * \chi_{K \cdot \varpi^{\mu_2^\vee} \cdot K} = \chi_{K \cdot \varpi^{\mu_1^\vee} \varpi^{\mu_2^\vee} \cdot K} = \chi_{K \cdot \varpi^{\mu_1^\vee \mu_2^\vee} \cdot K}.$$

Let R_K^+ be the algebra generated by the elements $\chi_{K \cdot \varpi^{\mu^\vee} \cdot K}$, where μ^\vee goes over all the dominant cocharacters. Then R_K^+ is a commutative subalgebra of $\mathcal{H}(G // K)$. Using the Cartan decomposition, we may decompose

$$\mathcal{H}(G // K) = \mathcal{H}(K // K) * R_K^+ * \mathcal{H}(K // K).$$

Here, $\mathcal{H}(K // K)$ is a finite-dimensional algebra, consisting of functions $K \rightarrow \mathbb{C}$ bi-invariant under K , and R_K^+ is abelian. This shows that $\mathcal{H}(K // K)$ breaks into two pieces: a small (finite-dimensional) non-commutative one, and a large (infinite-dimensional) abelian one.

Remark 3. An important feature that we will not discuss is the Jacquet functor. Suppose that $P = MU \subset G$ is a proper parabolic subgroup with Levi part M , and U^- is the radical opposite to U . Suppose that we have a Iwahori-type decomposition[1, Lemma 3.11]

$$K = (K \cap U^-) (K \cap M) (K \cap U).$$

(for instance, this holds for $K = \mathcal{K}$ and $K = \mathcal{I}$ from the next section). Let π be an admissible representation of G . Let

$$J_U(\pi) = \pi / \text{span}_{\mathbb{C}} \{ \pi(u)v - v \mid v \in \pi, u \in U \}.$$

Then the quotient map

$$\pi \rightarrow J_U(\pi)$$

defines a surjection for the subspaces of K -fixed vectors:

$$\pi^K \rightarrow J_U(\pi)^{K \cap M}.$$

This can be used to show that certain representations can be embedded as subrepresentations of principal series representations. See for example [4, Section 12].

2. IWAHORI–MATSUMOTO HECKE ALGEBRA

Consider the quotient map

$$\nu: \mathcal{K} \rightarrow G(\mathbb{F}_q).$$

The inverse image of $B(\mathbb{F}_q)$ under this map is called the *Iwahori subgroup* of $G(F)$. We denote it by \mathcal{I} . Assume henceforth that the Haar measure is normalized so that \mathcal{I} has measure 1.

Example 4. If $G = \text{GL}_n$ then

$$\mathcal{I} = \nu^{-1}(B(\mathbb{F}_q)) = \left\{ \begin{pmatrix} \mathfrak{o}^\times & \mathfrak{o} & \mathfrak{o} & \cdots & \mathfrak{o} \\ \mathfrak{p} & \mathfrak{o}^\times & \mathfrak{o} & \cdots & \mathfrak{o} \\ \mathfrak{p} & \mathfrak{p} & \ddots & \ddots & \vdots \\ \vdots & \vdots & \ddots & \mathfrak{o}^\times & \mathfrak{o} \\ \mathfrak{p} & \mathfrak{p} & \cdots & \mathfrak{p} & \mathfrak{o}^\times \end{pmatrix} \right\}.$$

Let $\tilde{W} = W \rtimes A(\varpi)$ be the extended affine Weyl group. We have the following Bruhat–Iwahori decomposition:

$$G = \bigsqcup_{x \in \tilde{W}} \mathcal{I} \cdot x \cdot \mathcal{I}.$$

It follows that $\{ \chi_{\mathcal{I} \cdot x \cdot \mathcal{I}} \}_{x \in \tilde{W}}$ forms a basis for $\mathcal{H}(G // \mathcal{I})$.

The group \tilde{W} is almost a Coxeter group. It has a well-known standard presentation. In general,

$$\tilde{W} = W_{\text{aff}} \rtimes \pi_1(G),$$

where $\pi_1(G)$ is the fundamental group of G and W_{aff} is the affine Weyl group, generated by all affine reflections. W_{aff} has a standard presentation as a Coxeter group.

Example 5. When $G = \text{GL}_n$, we have that W_{aff} is generated by s_1, \dots, s_n satisfying the relations

- (1) $s_j^2 = 1$.
- (2) $s_j s_k = s_k s_j$ if $j - k \not\equiv 0, \pm 1 \pmod{n}$.
- (3) $s_j s_{j+1} s_j = s_{j+1} s_j s_{j+1}$, where $j + 1$ is taken modulo n .

We have that $\pi_1(\mathrm{GL}_n) \cong \mathbb{Z}$ and that if $h \in \pi_1(\mathrm{GL}_n)$ is a generator, then for any j

$$hs_jh^{-1} = s_{j+1},$$

where $j+1$ is taken modulo n .

We may choose the following matrices to represent these elements: for $1 \leq j \leq n-1$, choose for s_j the permutation matrix that swaps the columns at positions j and $j+1$. Choose

$$s_n = \begin{pmatrix} & & \varpi^{-1} \\ & I_{n-2} & \\ \varpi & & \end{pmatrix}$$

and

$$h = \begin{pmatrix} & & \varpi^{-1} \\ 1 & & \\ & \ddots & \\ & & 1 \end{pmatrix}.$$

There is a notion of the length of an element in \tilde{W} : since W_{aff} is a Coxeter group, it comes with a length function $\ell: W_{\mathrm{aff}} \rightarrow \mathbb{Z}_{\geq 0}$ and we denote by $\ell: \tilde{W} \rightarrow W_{\mathrm{aff}} \rightarrow \mathbb{Z}_{\geq 0}$ the length function of \tilde{W} .

Example 6. If $G = \mathrm{GL}_n$ then any element in \tilde{W} can be written in the form

$$w = h^m \cdot s_{i_1} \cdot s_{i_2} \cdots s_{i_r},$$

where $m \in \mathbb{Z}$. We denote by $\ell(w)$ the minimal r such that w can be written in such form. Such presentation of w is called a *reduced expression*.

The elements of $\mathcal{H}(G // \mathcal{I})$ corresponding to the generators of \tilde{W} generate $\mathcal{H}(G // \mathcal{I})$ as an algebra. They satisfy the corresponding Hecke algebra relations.

Example 7. If $G = \mathrm{GL}_n$, let us define for any $w \in \tilde{W}$

$$f_w = \chi_{\mathcal{I} \cdot w \cdot \mathcal{I}}.$$

Then we have the following relations:

(1) $f_{s_j} * f_{s_j} = (q-1)f_{s_j} + q$. This can also be written as

$$(f_{s_j} - q) * (f_{s_j} + 1) = 0.$$

(2) $f_{s_j} * f_{s_k} = f_{s_k} * f_{s_j}$ if $j - k \not\equiv 0, \pm 1 \pmod{n}$.

(3) $f_{s_j} * f_{s_{j+1}} * f_{s_j} = f_{s_{j+1}} * f_{s_j} * f_{s_{j+1}}$, where $j+1$ is taken modulo n .

(4) $f_{h^{-1}} = f_h^{-1}$.

(5) $f_h * f_{s_j} * f_h^{-1} = f_{s_{j+1}}$, where $j+1$ is taken modulo n .

Remark 8. We may attach to W_{aff} an Iwahori–Matsumoto Hecke algebra $\mathcal{H}_v(W_{\mathrm{aff}})$. It is an algebra over $\mathbb{Z}[v, v^{-1}]$. The Iwahori–Matsumoto Hecke algebra is generated by the generators T_s for any generator s of W_{aff} and is subject to their relations, where we modify the quadratic relations $s^2 = 1$ to be

$$(T_s - v)(T_s + v^{-1}) = 0,$$

for every quadratic generator s of W_{aff} . Then Iwahori and Matsumoto proved that if G is semisimple then $\mathcal{H}(G // \mathcal{I})$ is isomorphic to $\mathcal{H}_{q^{\frac{1}{2}}}(W_{\mathrm{aff}})$. More generally, we have that

$$\mathcal{H}(\mathcal{K} // \mathcal{I}) \cong \mathcal{H}_{q^{\frac{1}{2}}}(W_{\mathrm{aff}})$$

and that

$$\mathcal{H}(G // \mathcal{I}) \cong \mathcal{H}(\mathcal{K} // \mathcal{I}) \otimes_{\mathbb{Z}[q^{\frac{1}{2}}, q^{-\frac{1}{2}}]} R_{\mathcal{I}},$$

by the isomorphism sending

$$\chi_{\mathcal{I} \cdot w \cdot \mathcal{I}} \otimes f \mapsto \chi_{\mathcal{I} \cdot w \cdot \mathcal{I}} * f.$$

See the definition of $R_{\mathcal{I}}$ below.

2.1. Bernstein–Zelevinsky relation. Denote for $w \in \tilde{W}$,

$$f_w = \chi_{\mathcal{I} \cdot w \cdot \mathcal{I}}.$$

Recall that $R_{\mathcal{I}}$ is the subalgebra of $\mathcal{H}(G // \mathcal{I})$ generated by $f_{\varpi^{\mu^\vee}}$ for every dominant cocharacter μ^\vee . If μ_1^\vee and μ_2^\vee are dominant cocharacters then we have that

$$\ell\left(\varpi^{\mu_1^\vee \mu_2^\vee}\right) = \ell\left(\varpi^{\mu_1^\vee} \varpi^{\mu_2^\vee}\right) = \ell\left(\varpi^{\mu_1^\vee}\right) + \ell\left(\varpi^{\mu_2^\vee}\right),$$

and

$$(2.1) \quad f_{\varpi^{\mu_1^\vee}} * f_{\varpi^{\mu_2^\vee}} = f_{\varpi^{\mu_1^\vee} \varpi^{\mu_2^\vee}} = f_{\varpi^{\mu_1^\vee \mu_2^\vee}}.$$

In particular $R_{\mathcal{I}}^+$ is commutative. It can be shown from the relations of the generators of the Hecke algebra that f_w is invertible for any $w \in \tilde{W}$. Let $R_{\mathcal{I}}$ be the algebra generated by $f_{\varpi^{\mu^\vee}}$ and $f_{\varpi^{\mu^\vee}}^{-1}$ for every dominant cocharacter μ^\vee . It seems tempting to define a map

$$A(\varpi) \rightarrow R_{\mathcal{I}}^\times$$

by the formula

$$\varpi^{\mu^\vee} \mapsto f_{\varpi^{\mu^\vee}}.$$

However, this map will not be a group homomorphism $A(\varpi) \rightarrow R_{\mathcal{I}}^\times$. This can be fixed as follows. First let us define a normalization of f_w :

$$T_w = q^{-\frac{\ell(w)}{2}} f_w.$$

If λ^\vee is a cocharacter, we can write $\lambda^\vee = \mu_1^\vee \cdot (\mu_2^\vee)^{-1}$ where μ_1^\vee and μ_2^\vee are dominant cocharacters. We define a map

$$\begin{aligned} \theta: A(\varpi) &\rightarrow R_{\mathcal{I}}^\times, \\ \theta(\lambda^\vee) &= T_{\mu_1^\vee} * (T_{\mu_2^\vee})^{-1}. \end{aligned}$$

This is well defined because of (2.1). This map is now a homomorphism.

Example 9. If $G = \mathrm{GL}_n$ then if we denote

$$b_j = \begin{pmatrix} I_{j-1} & & \\ & \varpi & \\ & & I_{n-j} \end{pmatrix},$$

then

$$b_j = a_j \cdot a_{j-1}^{-1},$$

and

$$\theta(b_j) = T_{a_j} * (T_{a_{j-1}})^{-1}.$$

One now needs to express a_{j-1} as a product of h^{-j} and of the generators s_1, \dots, s_{n-1} in order to be able to write an expression for $(T_{a_{j-1}})^{-1}$ as a product of the inverses of the

corresponding generators. In turn, these inverses are computed using the quadratic relations of the Hecke algebra.

Theorem 10 (Bernstein–Zelevinsky presentation). *Let s be any one of the generators of W_{aff} (as a Coxeter group). For any cocharacter λ^\vee we have that $\theta(\lambda^\vee) - \theta(s(\lambda^\vee))$ is divisible by $1 - \theta(\alpha^\vee)^{-1}$ in the ring $R_{\mathcal{I}}$ and the following equality holds:*

$$\begin{aligned} \theta(\lambda^\vee) T_s - T_s \theta(s(\lambda^\vee)) &= T_s \theta(\lambda^\vee) - \theta(s(\lambda^\vee)) T_s \\ &= \left(q^{\frac{1}{2}} - q^{-\frac{1}{2}} \right) \frac{\theta(\lambda^\vee) - \theta(s(\lambda^\vee))}{1 - \theta(\alpha^\vee)^{-1}}, \end{aligned}$$

where α^\vee is a certain fundamental cocharacter on which s acts by $s(\alpha^\vee) = (\alpha^\vee)^{-1}$.

Example 11. When $G = \text{GL}_n$, recall that λ^\vee is parameterized by $\lambda^\vee = \mu_{(k_1, \dots, k_n)}^\vee$ where $k_1, \dots, k_n \in \mathbb{Z}$. We have that $W = \langle s_i \mid 1 \leq i \leq n \rangle \cong S_n$ acts on λ^\vee by permuting the coordinates $(k_1, \dots, k_n) \in \mathbb{Z}$. In this case, $s = s_k$ for some k and

$$\alpha^\vee = \alpha_k^\vee = (0, 0, \dots, 0, -1, 1, 0, \dots, 0)$$

is the cocharacter corresponding to the matrix

$$\begin{pmatrix} I_{k-1} & & & \\ & \varpi^{-1} & & \\ & & \varpi & \\ & & & I_{n-k-1} \end{pmatrix},$$

so $\theta(\alpha^\vee) = T_{a_{k-1}} * T_{a_{k+1}} * T_{a_k}^{-2}$.

2.2. Center of $\mathcal{H}(G // \mathcal{I})$. From the Bernstein–Zelevinsky relation, the following description of the center of $\mathcal{H}(G // \mathcal{I})$ can be concluded.

Theorem 12. *The center of $\mathcal{H}(G // \mathcal{I})$ is the subspace of $R_{\mathcal{I}}^+$ consisting of elements of the form*

$$\sum_{\mu^\vee} a_{\mu^\vee} \theta(\mu^\vee)$$

such that $a_{w\mu^\vee} = a_{\mu^\vee}$ for every $w \in W$ and cocharacter μ^\vee . In particular, this center is isomorphic to $\mathbb{C}[A(\varpi)]^W$.

REFERENCES

- [1] I. N. Bernstein and A. V. Zelevinsky. Representations of the group $GL(n, F)$, where F is a local non-Archimedean field. *Uspehi Mat. Nauk*, 31(3(189)):5–70, 1976. 2
- [2] Daniel Bump. Hecke Algebras. Available at <http://sporadic.stanford.edu/bump/math263/hecke.pdf>.
- [3] Thomas J. Haines, Robert E. Kottwitz, and Amritanshu Prasad. Iwahori-Hecke algebras. *J. Ramanujan Math. Soc.*, 25(2):113–145, 2010.
- [4] Roger Howe. Hecke algebras and p -adic GL_n . In *Representation theory and analysis on homogeneous spaces (New Brunswick, NJ, 1993)*, volume 177 of *Contemp. Math.*, pages 65–100. Amer. Math. Soc., Providence, RI, 1994. 3
- [5] Roger Howe. Affine-like Hecke algebras and p -adic representation theory. In *Iwahori-Hecke algebras and their representation theory (Martina-Franca, 1999)*, volume 1804 of *Lecture Notes in Math.*, pages 27–69. Springer, Berlin, 2002.